

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Expected	32.4%
	Response at Six Months	Expected	11.8%
	Remission at Six Months	Expected	5.9%
	Follow-up at 12 Months	Expected	41.2%
	Response at 12 Months	Expected	23.5%
	Remission at 12 Months	Expected	11.8%
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Below Expected	38.0%
	Response at Six Months	Below Expected	12.5%
	Remission at Six Months	Below Expected	4.8%
	Follow-up at 12 Months	Below Expected	33.7%
	Response at 12 Months	Below Expected	11.8%
	Remission at 12 Months	Below Expected	6.3%
Adolescent Mental Health and/or Depression Screening*		Above	98.1%
Colorectal Cancer Screening		Above Expected	79.5%
Optimal Asthma Control - Adults		Above Expected	63.1%
Optimal Asthma Control - Children		Expected	61.8%
Optimal Diabetes Care		Above Expected	55.5%
HbA1c Control		Above Expected	84.0%
BP Control		Above Expected	84.8%
Daily Aspirin Use		Expected	98.5%
Statin Use		Expected	89.8%
Tobacco-Free		Expected	86.0%
Optimal Vascular Care		Expected	55.4%
BP Control		Above Expected	84.9%
Daily Aspirin Use		Below Expected	79.1%
Statin Use		Expected	92.1%
Tobacco-Free		Expected	83.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).