

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Below Expected	40.9%
	Response at Six Months	Below Expected	11.4%
	Remission at Six Months	Expected	5.4%
	Follow-up at 12 Months	Expected	52.4%
	Response at 12 Months	Expected	17.5%
	Remission at 12 Months	Expected	10.7%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Above Expected	89.0%
Optimal Asthma Control - Adults		Above Expected	74.4%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Above Expected	59.4%
HbA1c Control		Above Expected	84.7%
BP Control		Above Expected	86.6%
Daily Aspirin Use		Expected	97.6%
Statin Use		Expected	88.4%
Tobacco-Free		Expected	89.5%
Optimal Vascular Care		Expected	62.1%
BP Control		Expected	85.2%
Daily Aspirin Use		Expected	85.6%
Statin Use		Expected	91.7%
Tobacco-Free		Expected	88.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).