

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Above Expected	85.7%
Optimal Asthma Control - Adults		Expected	49.8%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	52.7%
HbA1c Control		Above Expected	82.1%
BP Control		Below Expected	72.5%
Daily Aspirin Use		Expected	98.3%
Statin Use		Expected	89.9%
Tobacco-Free		Above Expected	96.7%
Optimal Vascular Care		Below Expected	53.1%
BP Control		Below Expected	72.9%
Daily Aspirin Use		Below Expected	78.7%
Statin Use		Expected	94.3%
Tobacco-Free		Above Expected	96.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).