

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Above	95.5%
Colorectal Cancer Screening		Below Expected	69.4%
Optimal Asthma Control - Adults		Expected	52.1%
Optimal Asthma Control - Children		Expected	48.4%
Optimal Diabetes Care		Below Expected	45.1%
HbA1c Control		Below Expected	70.9%
BP Control		Above Expected	84.4%
Daily Aspirin Use		Above Expected	99.3%
Statin Use		Expected	86.7%
Tobacco-Free		Expected	85.2%
Optimal Vascular Care		Expected	60.9%
BP Control		Above Expected	86.3%
Daily Aspirin Use		Above Expected	90.7%
Statin Use		Expected	93.4%
Tobacco-Free		Expected	80.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).