

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	81.5%
	Follow-up at Six Months	Below Expected	25.3%
	Response at Six Months	Below Expected	9.3%
	Remission at Six Months	Below Expected	3.0%
	Follow-up at 12 Months	Below Expected	27.0%
	Response at 12 Months	Below Expected	11.4%
	Remission at 12 Months	Expected	7.6%
Adolescent Mental Health and/or Depression Screening*		Above	95.8%
Colorectal Cancer Screening		Below Expected	68.9%
Optimal Asthma Control - Adults		Below Expected	3.5%
Optimal Asthma Control - Children		Below Expected	2.2%
Optimal Diabetes Care		Below Expected	38.1%
HbA1c Control		Below Expected	70.1%
BP Control		Expected	78.7%
Daily Aspirin Use		Expected	98.0%
Statin Use		Below Expected	82.1%
Tobacco-Free		Expected	84.3%
Optimal Vascular Care		Below Expected	48.3%
BP Control		Expected	79.7%
Daily Aspirin Use		Below Expected	80.8%
Statin Use		Expected	90.1%
Tobacco-Free		Expected	81.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).