

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	79.8%
	Follow-up at Six Months	Below Expected	36.6%
	Response at Six Months	Below Expected	9.4%
	Remission at Six Months	Expected	5.9%
	Follow-up at 12 Months	Below Expected	34.2%
	Response at 12 Months	Expected	12.9%
	Remission at 12 Months	Expected	9.9%
Adolescent Mental Health and/or Depression Screening*		Above	100.0%
Colorectal Cancer Screening		Below Expected	67.3%
Optimal Asthma Control - Adults		Below Expected	2.2%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	46.0%
HbA1c Control		Expected	74.1%
BP Control		Above Expected	87.6%
Daily Aspirin Use		Expected	98.1%
Statin Use		Expected	88.0%
Tobacco-Free		Expected	81.9%
Optimal Vascular Care		Expected	50.6%
BP Control		Above Expected	90.0%
Daily Aspirin Use		Below Expected	80.0%
Statin Use		Expected	92.0%
Tobacco-Free		Below Expected	74.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).