

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	66.4%
	Follow-up at Six Months	Expected	34.0%
	Response at Six Months	Expected	14.9%
	Remission at Six Months	Expected	10.6%
	Follow-up at 12 Months	Expected	38.3%
	Response at 12 Months	Expected	10.6%
	Remission at 12 Months	Expected	8.5%
Adolescent Mental Health and/or Depression Screening*		Average	89.1%
Colorectal Cancer Screening		Expected	65.9%
Optimal Asthma Control - Adults		Expected	38.7%
Optimal Asthma Control - Children		Expected	33.3%
Optimal Diabetes Care		Expected	47.2%
HbA1c Control		Above Expected	79.9%
BP Control		Expected	87.0%
Daily Aspirin Use		Below Expected	94.0%
Statin Use		Expected	84.5%
Tobacco-Free		Expected	79.9%
Optimal Vascular Care		Expected	46.1%
BP Control		Expected	83.1%
Daily Aspirin Use		Below Expected	73.1%
Statin Use		Expected	87.7%
Tobacco-Free		Expected	82.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).