

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	76.0%
	Follow-up at Six Months	Below Expected	25.4%
	Response at Six Months	Expected	11.3%
	Remission at Six Months	Expected	9.9%
	Follow-up at 12 Months	Below Expected	25.4%
	Response at 12 Months	Expected	21.1%
	Remission at 12 Months	Expected	15.5%
Adolescent Mental Health and/or Depression Screening*		Below	60.5%
Colorectal Cancer Screening		Below Expected	61.7%
Optimal Asthma Control - Adults		Below Expected	0.0%
Optimal Asthma Control - Children		Below Expected	0.0%
Optimal Diabetes Care		Expected	42.1%
HbA1c Control		Expected	68.9%
BP Control		Expected	81.0%
Daily Aspirin Use		Expected	99.7%
Statin Use		Expected	90.6%
Tobacco-Free		Below Expected	76.1%
Optimal Vascular Care		Expected	47.2%
BP Control		Expected	78.2%
Daily Aspirin Use		Above Expected	93.3%
Statin Use		Expected	91.2%
Tobacco-Free		Below Expected	69.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).