

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	66.3%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Below	72.7%
Colorectal Cancer Screening		Below Expected	58.0%
Optimal Asthma Control - Adults		Expected	36.5%
Optimal Asthma Control - Children		Expected	45.7%
Optimal Diabetes Care		Expected	41.8%
HbA1c Control		Expected	72.5%
BP Control		Above Expected	88.3%
Daily Aspirin Use		Expected	96.4%
Statin Use		Expected	84.7%
Tobacco-Free		Expected	78.6%
Optimal Vascular Care		Below Expected	34.9%
BP Control		Expected	82.6%
Daily Aspirin Use		Expected	79.1%
Statin Use		Expected	90.7%
Tobacco-Free		Below Expected	65.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).