

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	83.0%
	Follow-up at Six Months	Below Expected	19.8%
	Response at Six Months	Below Expected	4.4%
	Remission at Six Months	Below Expected	2.2%
	Follow-up at 12 Months	Below Expected	29.7%
	Response at 12 Months	Below Expected	8.8%
	Remission at 12 Months	Expected	4.4%
Adolescent Mental Health and/or Depression Screening*		Average	92.1%
Colorectal Cancer Screening		Expected	68.0%
Optimal Asthma Control - Adults		Below Expected	5.6%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	39.0%
HbA1c Control		Expected	69.8%
BP Control		Expected	76.8%
Daily Aspirin Use		Expected	98.3%
Statin Use		Expected	87.5%
Tobacco-Free		Expected	86.2%
Optimal Vascular Care		Expected	45.2%
BP Control		Expected	75.7%
Daily Aspirin Use		Expected	79.5%
Statin Use		Expected	89.5%
Tobacco-Free		Expected	80.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).