

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	25.8%
	Follow-up at Six Months	Below Expected	12.7%
	Response at Six Months	Expected	5.5%
	Remission at Six Months	Below Expected	0.0%
	Follow-up at 12 Months	Expected	32.7%
	Response at 12 Months	Expected	7.3%
	Remission at 12 Months	Expected	1.8%
Adolescent Mental Health and/or Depression Screening*		Below	58.7%
Colorectal Cancer Screening		Above Expected	70.1%
Optimal Asthma Control - Adults		Below Expected	30.9%
Optimal Asthma Control - Children		Below Expected	33.3%
Optimal Diabetes Care		Expected	39.7%
HbA1c Control		Expected	68.1%
BP Control		Above Expected	87.8%
Daily Aspirin Use		Below Expected	96.1%
Statin Use		Expected	82.4%
Tobacco-Free		Expected	82.5%
Optimal Vascular Care		Expected	50.6%
BP Control		Above Expected	90.3%
Daily Aspirin Use		Below Expected	76.0%
Statin Use		Below Expected	85.9%
Tobacco-Free		Expected	82.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).