

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	86.9%
	Follow-up at Six Months	Below Expected	26.9%
	Response at Six Months	Below Expected	6.0%
	Remission at Six Months	Below Expected	2.5%
	Follow-up at 12 Months	Below Expected	28.9%
	Response at 12 Months	Below Expected	11.0%
	Remission at 12 Months	Expected	6.5%
Adolescent Mental Health and/or Depression Screening*		Above	98.4%
Colorectal Cancer Screening		Below Expected	40.7%
Optimal Asthma Control - Adults		Below Expected	13.4%
Optimal Asthma Control - Children		Below Expected	8.3%
Optimal Diabetes Care		Expected	43.4%
HbA1c Control		Expected	73.4%
BP Control		Expected	81.0%
Daily Aspirin Use		Expected	97.9%
Statin Use		Expected	87.5%
Tobacco-Free		Below Expected	81.7%
Optimal Vascular Care		Expected	51.9%
BP Control		Expected	81.7%
Daily Aspirin Use		Expected	83.2%
Statin Use		Expected	88.2%
Tobacco-Free		Expected	82.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).