

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	80.8%
	Follow-up at Six Months	Below Expected	34.0%
	Response at Six Months	Expected	11.4%
	Remission at Six Months	Expected	7.8%
	Follow-up at 12 Months	Below Expected	36.9%
	Response at 12 Months	Expected	13.5%
	Remission at 12 Months	Expected	8.5%
Adolescent Mental Health and/or Depression Screening*		Average	96.2%
Colorectal Cancer Screening		Expected	72.1%
Optimal Asthma Control - Adults		Below Expected	7.7%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	46.7%
HbA1c Control		Expected	73.7%
BP Control		Expected	85.7%
Daily Aspirin Use		Expected	98.0%
Statin Use		Expected	85.9%
Tobacco-Free		Expected	86.3%
Optimal Vascular Care		Expected	52.4%
BP Control		Expected	84.0%
Daily Aspirin Use		Below Expected	77.8%
Statin Use		Expected	91.5%
Tobacco-Free		Expected	84.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).