

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	96.3%
	Follow-up at Six Months	Expected	66.7%
	Response at Six Months	Expected	26.4%
	Remission at Six Months	Expected	15.3%
	Follow-up at 12 Months	Expected	63.9%
	Response at 12 Months	Expected	29.2%
	Remission at 12 Months	Expected	18.1%
Adolescent Mental Health and/or Depression Screening*		Average	91.8%
Colorectal Cancer Screening		Expected	69.2%
Optimal Asthma Control - Adults		Expected	65.2%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	49.9%
HbA1c Control		Expected	77.4%
BP Control		Expected	79.0%
Daily Aspirin Use		Expected	99.7%
Statin Use		Expected	91.7%
Tobacco-Free		Expected	84.7%
Optimal Vascular Care		Expected	57.9%
BP Control		Expected	80.2%
Daily Aspirin Use		Expected	89.9%
Statin Use		Above Expected	97.5%
Tobacco-Free		Expected	82.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).