

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Expected	67.8%
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
<b>Optimal Diabetes Care</b>		<b>Expected</b>	<b>39.1%</b>
HbA1c Control		Expected	71.0%
BP Control		Expected	76.8%
Daily Aspirin Use		Expected	97.1%
Statin Use		Expected	92.8%
Tobacco-Free		Expected	81.2%
<b>Optimal Vascular Care</b>		<b>Expected</b>	<b>48.8%</b>
BP Control		Expected	70.7%
Daily Aspirin Use		Expected	85.4%
Statin Use		Expected	80.5%
Tobacco-Free		Expected	78.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).