

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	64.2%
	Follow-up at Six Months	Below Expected	29.4%
	Response at Six Months	Below Expected	10.3%
	Remission at Six Months	Expected	5.2%
	Follow-up at 12 Months	Below Expected	28.9%
	Response at 12 Months	Below Expected	10.3%
	Remission at 12 Months	Expected	5.2%
Adolescent Mental Health and/or Depression Screening*		Average	91.1%
Colorectal Cancer Screening		Below Expected	66.3%
Optimal Asthma Control - Adults		Expected	40.9%
Optimal Asthma Control - Children		Expected	38.4%
Optimal Diabetes Care		Expected	43.0%
HbA1c Control		Expected	71.5%
BP Control		Above Expected	89.3%
Daily Aspirin Use		Expected	97.9%
Statin Use		Below Expected	80.3%
Tobacco-Free		Below Expected	81.1%
Optimal Vascular Care		Expected	53.3%
BP Control		Above Expected	86.5%
Daily Aspirin Use		Expected	81.7%
Statin Use		Below Expected	85.5%
Tobacco-Free		Expected	79.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).