

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	37.4%
	Response at Six Months	Expected	10.8%
	Remission at Six Months	Expected	6.0%
	Follow-up at 12 Months	Below Expected	24.1%
	Response at 12 Months	Expected	10.8%
	Remission at 12 Months	Expected	7.2%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Below Expected	33.2%
Optimal Asthma Control - Adults		Below Expected	22.5%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	12.1%
HbA1c Control		Expected	66.7%
BP Control		Expected	72.7%
Daily Aspirin Use		Expected	97.0%
Statin Use		Expected	87.9%
Tobacco-Free		Below Expected	37.9%
Optimal Vascular Care		Below Expected	17.7%
BP Control		Expected	70.6%
Daily Aspirin Use		Expected	70.6%
Statin Use		Expected	76.5%
Tobacco-Free		Below Expected	38.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).