

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	45.6%
	Follow-up at Six Months	Expected	37.2%
	Response at Six Months	Expected	15.0%
	Remission at Six Months	Expected	5.3%
	Follow-up at 12 Months	Expected	42.5%
	Response at 12 Months	Expected	15.9%
	Remission at 12 Months	Expected	9.7%
Adolescent Mental Health and/or Depression Screening*		Below	64.7%
Colorectal Cancer Screening		Above Expected	73.7%
Optimal Asthma Control - Adults		Below Expected	26.8%
Optimal Asthma Control - Children		Below Expected	9.3%
Optimal Diabetes Care		Expected	48.9%
HbA1c Control		Above Expected	79.7%
BP Control		Above Expected	88.8%
Daily Aspirin Use		Below Expected	96.2%
Statin Use		Expected	83.9%
Tobacco-Free		Below Expected	79.1%
Optimal Vascular Care		Below Expected	42.6%
BP Control		Above Expected	86.9%
Daily Aspirin Use		Below Expected	74.6%
Statin Use		Expected	87.5%
Tobacco-Free		Below Expected	75.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).