

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	97.7%
	Follow-up at Six Months	Expected	37.3%
	Response at Six Months	Expected	9.0%
	Remission at Six Months	Expected	7.5%
	Follow-up at 12 Months	Expected	50.8%
	Response at 12 Months	Expected	14.9%
	Remission at 12 Months	Expected	9.0%
Adolescent Mental Health and/or Depression Screening*		Average	93.2%
Colorectal Cancer Screening		Below Expected	56.7%
Optimal Asthma Control - Adults		Expected	43.3%
Optimal Asthma Control - Children		Below Expected	23.5%
Optimal Diabetes Care		Below Expected	33.6%
HbA1c Control		Expected	69.2%
BP Control		Below Expected	70.6%
Daily Aspirin Use		Expected	99.3%
Statin Use		Below Expected	77.6%
Tobacco-Free		Expected	80.4%
Optimal Vascular Care		Expected	53.3%
BP Control		Expected	79.2%
Daily Aspirin Use		Expected	83.1%
Statin Use		Expected	88.3%
Tobacco-Free		Expected	80.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).