

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	83.4%
	Follow-up at Six Months	Expected	46.2%
	Response at Six Months	Expected	17.4%
	Remission at Six Months	Expected	9.7%
	Follow-up at 12 Months	Expected	50.5%
	Response at 12 Months	Expected	21.4%
	Remission at 12 Months	Expected	12.7%
Adolescent Mental Health and/or Depression Screening*		Above	100.0%
Colorectal Cancer Screening		Above Expected	80.4%
Optimal Asthma Control - Adults		Above Expected	64.3%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Above Expected	52.8%
HbA1c Control		Expected	76.2%
BP Control		Expected	81.4%
Daily Aspirin Use		Expected	98.5%
Statin Use		Expected	91.2%
Tobacco-Free		Above Expected	88.1%
Optimal Vascular Care		Expected	61.1%
BP Control		Expected	83.7%
Daily Aspirin Use		Expected	83.7%
Statin Use		Expected	94.7%
Tobacco-Free		Above Expected	89.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).