

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	88.1%
	Follow-up at Six Months	Below Expected	43.1%
	Response at Six Months	Expected	14.0%
	Remission at Six Months	Below Expected	5.7%
	Follow-up at 12 Months	Expected	43.9%
	Response at 12 Months	Expected	14.8%
	Remission at 12 Months	Expected	7.8%
Adolescent Mental Health and/or Depression Screening*		Above	100.0%
Colorectal Cancer Screening		Above Expected	75.6%
Optimal Asthma Control - Adults		Above Expected	69.0%
Optimal Asthma Control - Children		Expected	71.4%
Optimal Diabetes Care		Above Expected	51.8%
HbA1c Control		Expected	76.5%
BP Control		Expected	83.2%
Daily Aspirin Use		Expected	97.9%
Statin Use		Above Expected	91.0%
Tobacco-Free		Expected	87.4%
Optimal Vascular Care		Expected	55.9%
BP Control		Expected	82.5%
Daily Aspirin Use		Expected	83.6%
Statin Use		Expected	93.7%
Tobacco-Free		Expected	83.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).