

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	89.4%
	Follow-up at Six Months	Expected	52.5%
	Response at Six Months	Expected	17.4%
	Remission at Six Months	Expected	10.1%
	Follow-up at 12 Months	Expected	50.3%
	Response at 12 Months	Expected	18.0%
	Remission at 12 Months	Expected	10.8%
Adolescent Mental Health and/or Depression Screening*		Above	100.0%
Colorectal Cancer Screening		Expected	64.6%
Optimal Asthma Control - Adults		Above Expected	63.0%
Optimal Asthma Control - Children		Above Expected	63.8%
Optimal Diabetes Care		Expected	39.1%
HbA1c Control		Below Expected	64.5%
BP Control		Expected	78.5%
Daily Aspirin Use		Expected	99.2%
Statin Use		Above Expected	91.1%
Tobacco-Free		Expected	81.8%
Optimal Vascular Care		Expected	50.9%
BP Control		Expected	76.5%
Daily Aspirin Use		Expected	90.5%
Statin Use		Above Expected	94.5%
Tobacco-Free		Expected	75.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).