

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	91.7%
	Follow-up at Six Months	Expected	46.4%
	Response at Six Months	Expected	17.5%
	Remission at Six Months	Expected	8.1%
	Follow-up at 12 Months	Expected	56.6%
	Response at 12 Months	Expected	22.1%
	Remission at 12 Months	Expected	12.3%
Adolescent Mental Health and/or Depression Screening*		Above	97.3%
Colorectal Cancer Screening		Above Expected	75.7%
Optimal Asthma Control - Adults		Above Expected	75.4%
Optimal Asthma Control - Children		Expected	68.9%
Optimal Diabetes Care		Above Expected	54.8%
HbA1c Control		Expected	79.3%
BP Control		Expected	83.8%
Daily Aspirin Use		Expected	98.4%
Statin Use		Expected	91.0%
Tobacco-Free		Expected	88.3%
Optimal Vascular Care		Expected	59.1%
BP Control		Expected	82.7%
Daily Aspirin Use		Expected	81.9%
Statin Use		Expected	91.1%
Tobacco-Free		Above Expected	92.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).