

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	78.4%
	Follow-up at Six Months	Below Expected	33.3%
	Response at Six Months	Below Expected	7.8%
	Remission at Six Months	Below Expected	3.3%
	Follow-up at 12 Months	Expected	35.6%
	Response at 12 Months	Expected	12.2%
	Remission at 12 Months	Expected	10.0%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Above Expected	78.4%
Optimal Asthma Control - Adults		Below Expected	39.6%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	43.2%
HbA1c Control		Expected	72.5%
BP Control		Below Expected	71.0%
Daily Aspirin Use		Expected	98.3%
Statin Use		Expected	89.2%
Tobacco-Free		Above Expected	88.4%
Optimal Vascular Care		Expected	53.2%
BP Control		Below Expected	75.1%
Daily Aspirin Use		Expected	86.1%
Statin Use		Expected	92.7%
Tobacco-Free		Above Expected	87.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).