

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Below Expected	49.4%
Optimal Asthma Control - Adults		Expected	46.0%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	32.4%
HbA1c Control		Below Expected	58.4%
BP Control		Below Expected	62.0%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	89.0%
Tobacco-Free		Expected	87.4%
Optimal Vascular Care		Expected	49.0%
BP Control		Below Expected	69.9%
Daily Aspirin Use		Expected	89.5%
Statin Use		Above Expected	96.7%
Tobacco-Free		Expected	84.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).