

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	68.8%
	Follow-up at Six Months	Above Expected	75.4%
	Response at Six Months	Expected	33.3%
	Remission at Six Months	Above Expected	28.1%
	Follow-up at 12 Months	Expected	56.1%
	Response at 12 Months	Expected	26.3%
	Remission at 12 Months	Expected	17.5%
Adolescent Mental Health and/or Depression Screening*		Average	95.2%
Colorectal Cancer Screening		Above Expected	74.2%
Optimal Asthma Control - Adults		Expected	52.9%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	42.8%
HbA1c Control		Expected	73.3%
BP Control		Expected	81.3%
Daily Aspirin Use		Expected	98.9%
Statin Use		Expected	93.6%
Tobacco-Free		Below Expected	77.0%
Optimal Vascular Care		Expected	52.0%
BP Control		Expected	81.6%
Daily Aspirin Use		Above Expected	94.2%
Statin Use		Expected	95.1%
Tobacco-Free		Below Expected	66.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).