

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	82.4%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	85.7%
	Follow-up at Six Months	Expected	56.8%
	Response at Six Months	Expected	15.6%
	Remission at Six Months	Expected	8.3%
	Follow-up at 12 Months	Above Expected	69.4%
	Response at 12 Months	Above Expected	24.7%
	Remission at 12 Months	Above Expected	16.4%
Adolescent Mental Health and/or Depression Screening*		Average	92.6%
Colorectal Cancer Screening		Above Expected	75.8%
Optimal Asthma Control - Adults		Above Expected	67.9%
Optimal Asthma Control - Children		Above Expected	65.9%
Optimal Diabetes Care		Expected	43.9%
HbA1c Control		Above Expected	75.5%
BP Control		Expected	82.6%
Daily Aspirin Use		Expected	98.6%
Statin Use		Expected	90.2%
Tobacco-Free		Below Expected	74.0%
Optimal Vascular Care		Expected	49.0%
BP Control		Expected	83.3%
Daily Aspirin Use		Expected	87.8%
Statin Use		Expected	93.5%
Tobacco-Free		Below Expected	69.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).