

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	91.7%
	Follow-up at Six Months	Above Expected	61.1%
	Response at Six Months	Expected	16.8%
	Remission at Six Months	Expected	9.4%
	Follow-up at 12 Months	Above Expected	62.6%
	Response at 12 Months	Expected	24.6%
	Remission at 12 Months	Expected	13.8%
Adolescent Mental Health and/or Depression Screening*		Average	89.2%
Colorectal Cancer Screening		Above Expected	79.4%
Optimal Asthma Control - Adults		Above Expected	66.5%
Optimal Asthma Control - Children		Expected	59.6%
Optimal Diabetes Care		Expected	46.8%
HbA1c Control		Expected	76.6%
BP Control		Expected	83.2%
Daily Aspirin Use		Expected	98.0%
Statin Use		Expected	88.5%
Tobacco-Free		Below Expected	77.2%
Optimal Vascular Care		Expected	53.9%
BP Control		Expected	84.4%
Daily Aspirin Use		Expected	87.1%
Statin Use		Expected	93.8%
Tobacco-Free		Below Expected	72.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).