

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	89.4%
	Follow-up at Six Months	Expected	42.3%
	Response at Six Months	Expected	15.5%
	Remission at Six Months	Expected	9.9%
	Follow-up at 12 Months	Expected	56.3%
	Response at 12 Months	Expected	26.8%
	Remission at 12 Months	Expected	15.5%
Adolescent Mental Health and/or Depression Screening*		Below	83.1%
Colorectal Cancer Screening		Above Expected	74.5%
Optimal Asthma Control - Adults		Above Expected	67.3%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	38.2%
HbA1c Control		Expected	75.2%
BP Control		Expected	84.7%
Daily Aspirin Use		Expected	98.0%
Statin Use		Below Expected	80.7%
Tobacco-Free		Below Expected	75.6%
Optimal Vascular Care		Expected	52.6%
BP Control		Expected	88.3%
Daily Aspirin Use		Expected	83.2%
Statin Use		Expected	89.8%
Tobacco-Free		Below Expected	70.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).