

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	72.5%
	Follow-up at Six Months	Expected	35.5%
	Response at Six Months	Expected	12.2%
	Remission at Six Months	Expected	8.4%
	Follow-up at 12 Months	Below Expected	20.6%
	Response at 12 Months	Below Expected	3.7%
	Remission at 12 Months	Below Expected	0.9%
Adolescent Mental Health and/or Depression Screening*		Average	90.6%
Colorectal Cancer Screening		Below Expected	53.5%
Optimal Asthma Control - Adults		Expected	41.5%
Optimal Asthma Control - Children		Expected	60.0%
Optimal Diabetes Care		Expected	33.9%
HbA1c Control		Expected	62.2%
BP Control		Below Expected	74.4%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	88.0%
Tobacco-Free		Expected	82.7%
Optimal Vascular Care		Expected	40.4%
BP Control		Below Expected	67.9%
Daily Aspirin Use		Expected	82.6%
Statin Use		Expected	90.8%
Tobacco-Free		Expected	76.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).