

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	90.9%
	Follow-up at Six Months	Above Expected	66.7%
	Response at Six Months	Expected	26.2%
	Remission at Six Months	Expected	11.9%
	Follow-up at 12 Months	Above Expected	61.9%
	Response at 12 Months	Expected	26.2%
	Remission at 12 Months	Expected	14.3%
Adult Depression	PHQ-9 Utilization*	Below	74.1%
	Follow-up at Six Months	Below Expected	26.9%
	Response at Six Months	Below Expected	9.8%
	Remission at Six Months	Expected	6.3%
	Follow-up at 12 Months	Below Expected	28.7%
	Response at 12 Months	Below Expected	9.4%
	Remission at 12 Months	Expected	5.6%
Adolescent Mental Health and/or Depression Screening*		Below	72.2%
Colorectal Cancer Screening		Below Expected	49.4%
Optimal Asthma Control - Adults		Expected	42.7%
Optimal Asthma Control - Children		Expected	46.8%
Optimal Diabetes Care		Below Expected	25.2%
HbA1c Control		Below Expected	55.7%
BP Control		Below Expected	70.9%
Daily Aspirin Use		Expected	98.7%
Statin Use		Below Expected	81.1%
Tobacco-Free		Expected	80.6%
Optimal Vascular Care		Below Expected	29.8%
BP Control		Below Expected	63.3%
Daily Aspirin Use		Below Expected	75.4%
Statin Use		Expected	87.1%
Tobacco-Free		Expected	70.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).