

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	60.6%
	Response at Six Months	Expected	24.0%
	Remission at Six Months	Expected	11.4%
	Follow-up at 12 Months	Expected	56.0%
	Response at 12 Months	Expected	24.6%
	Remission at 12 Months	Expected	16.0%
Adolescent Mental Health and/or Depression Screening*		Below	58.2%
Colorectal Cancer Screening		Above Expected	82.1%
Optimal Asthma Control - Adults		Expected	47.4%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Above Expected	69.8%
HbA1c Control		Above Expected	84.4%
BP Control		Above Expected	92.4%
Daily Aspirin Use		Expected	99.5%
Statin Use		Above Expected	93.8%
Tobacco-Free		Above Expected	93.1%
Optimal Vascular Care		Above Expected	70.3%
BP Control		Above Expected	91.3%
Daily Aspirin Use		Expected	87.3%
Statin Use		Expected	93.9%
Tobacco-Free		Expected	87.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).