

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Below	61.8%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	32.8%
	Follow-up at Six Months	Below Expected	27.4%
	Response at Six Months	Below Expected	9.1%
	Remission at Six Months	Expected	4.6%
	Follow-up at 12 Months	Below Expected	21.5%
	Response at 12 Months	Below Expected	7.3%
	Remission at 12 Months	Below Expected	3.7%
Adolescent Mental Health and/or Depression Screening*		Below	86.4%
Colorectal Cancer Screening		Below Expected	31.6%
Optimal Asthma Control - Adults		Below Expected	4.4%
Optimal Asthma Control - Children		Below Expected	14.9%
Optimal Diabetes Care		Below Expected	27.4%
HbA1c Control		Below Expected	60.3%
BP Control		Below Expected	66.5%
Daily Aspirin Use		Above Expected	99.8%
Statin Use		Expected	85.8%
Tobacco-Free		Expected	73.7%
Optimal Vascular Care		Expected	42.2%
BP Control		Expected	67.7%
Daily Aspirin Use		Expected	90.2%
Statin Use		Expected	94.1%
Tobacco-Free		Expected	65.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).