

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	78.8%
	Follow-up at Six Months	Expected	38.2%
	Response at Six Months	Expected	11.8%
	Remission at Six Months	Expected	2.9%
	Follow-up at 12 Months	Expected	41.2%
	Response at 12 Months	Expected	14.7%
	Remission at 12 Months	Expected	0.0%
Adult Depression	PHQ-9 Utilization*	Below	57.6%
	Follow-up at Six Months	Below Expected	34.7%
	Response at Six Months	Expected	11.4%
	Remission at Six Months	Expected	4.5%
	Follow-up at 12 Months	Expected	37.1%
	Response at 12 Months	Expected	13.1%
	Remission at 12 Months	Expected	6.5%
Adolescent Mental Health and/or Depression Screening*		Below	82.7%
Colorectal Cancer Screening		Above Expected	75.9%
Optimal Asthma Control - Adults		Below Expected	20.7%
Optimal Asthma Control - Children		Below Expected	22.7%
Optimal Diabetes Care		Below Expected	34.7%
HbA1c Control		Expected	66.5%
BP Control		Expected	82.5%
Daily Aspirin Use		Below Expected	97.1%
Statin Use		Below Expected	82.2%
Tobacco-Free		Below Expected	75.9%
Optimal Vascular Care		Below Expected	40.0%
BP Control		Expected	81.6%
Daily Aspirin Use		Below Expected	74.5%
Statin Use		Below Expected	85.6%
Tobacco-Free		Expected	75.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).