

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	40.2%
	Follow-up at Six Months	Below Expected	33.9%
	Response at Six Months	Expected	18.6%
	Remission at Six Months	Expected	8.9%
	Follow-up at 12 Months	Below Expected	33.1%
	Response at 12 Months	Expected	15.3%
	Remission at 12 Months	Expected	9.7%
Adolescent Mental Health and/or Depression Screening*		Below	31.3%
Colorectal Cancer Screening		Below Expected	69.8%
Optimal Asthma Control - Adults		Expected	55.2%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	49.3%
HbA1c Control		Expected	81.5%
BP Control		Below Expected	72.7%
Daily Aspirin Use		Expected	97.8%
Statin Use		Expected	88.1%
Tobacco-Free		Above Expected	95.6%
Optimal Vascular Care		Below Expected	33.3%
BP Control		Below Expected	63.7%
Daily Aspirin Use		Below Expected	63.1%
Statin Use		Expected	89.3%
Tobacco-Free		Expected	88.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).