

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	90.5%
	Follow-up at Six Months	Expected	53.7%
	Response at Six Months	Expected	29.6%
	Remission at Six Months	Expected	16.7%
	Follow-up at 12 Months	Expected	48.2%
	Response at 12 Months	Expected	20.4%
	Remission at 12 Months	Expected	14.8%
Adult Depression	PHQ-9 Utilization*	Above	90.1%
	Follow-up at Six Months	Above Expected	65.6%
	Response at Six Months	Above Expected	30.8%
	Remission at Six Months	Above Expected	18.7%
	Follow-up at 12 Months	Above Expected	67.0%
	Response at 12 Months	Expected	25.3%
	Remission at 12 Months	Expected	13.6%
Adolescent Mental Health and/or Depression Screening*		Above	98.0%
Colorectal Cancer Screening		Above Expected	80.0%
Optimal Asthma Control - Adults		Above Expected	72.3%
Optimal Asthma Control - Children		Above Expected	72.5%
Optimal Diabetes Care		Expected	55.0%
HbA1c Control		Expected	77.9%
BP Control		Expected	84.5%
Daily Aspirin Use		Expected	99.0%
Statin Use		Above Expected	93.1%
Tobacco-Free		Expected	85.9%
Optimal Vascular Care		Expected	54.2%
BP Control		Expected	84.9%
Daily Aspirin Use		Expected	86.2%
Statin Use		Expected	90.6%
Tobacco-Free		Expected	79.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).