

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Average	94.9%
Colorectal Cancer Screening		Below Expected	67.8%
Optimal Asthma Control - Adults		Expected	47.9%
Optimal Asthma Control - Children		Expected	59.7%
Optimal Diabetes Care		Above Expected	58.9%
HbA1c Control		Expected	80.8%
BP Control		Expected	86.9%
Daily Aspirin Use		Above Expected	100.0%
Statin Use		Expected	90.0%
Tobacco-Free		Expected	89.1%
Optimal Vascular Care		Expected	64.8%
BP Control		Expected	86.9%
Daily Aspirin Use		Above Expected	93.0%
Statin Use		Expected	93.0%
Tobacco-Free		Expected	83.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).