

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Expected	56.7%
	Response at Six Months	Expected	10.0%
	Remission at Six Months	Expected	3.3%
	Follow-up at 12 Months	Expected	40.0%
	Response at 12 Months	Expected	10.0%
	Remission at 12 Months	Expected	3.3%
Adult Depression	PHQ-9 Utilization*	Above	96.6%
	Follow-up at Six Months	Above Expected	72.1%
	Response at Six Months	Above Expected	27.0%
	Remission at Six Months	Above Expected	15.4%
	Follow-up at 12 Months	Above Expected	64.3%
	Response at 12 Months	Expected	23.6%
	Remission at 12 Months	Expected	14.1%
Adolescent Mental Health and/or Depression Screening*		Above	97.1%
Colorectal Cancer Screening		Below Expected	56.5%
Optimal Asthma Control - Adults		Expected	47.6%
Optimal Asthma Control - Children		Expected	50.2%
Optimal Diabetes Care		Below Expected	40.2%
HbA1c Control		Expected	73.4%
BP Control		Below Expected	73.1%
Daily Aspirin Use		Expected	98.2%
Statin Use		Expected	88.5%
Tobacco-Free		Expected	82.6%
Optimal Vascular Care		Below Expected	44.9%
BP Control		Below Expected	75.1%
Daily Aspirin Use		Expected	83.1%
Statin Use		Below Expected	87.3%
Tobacco-Free		Expected	79.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).