

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	89.5%
	Follow-up at Six Months	Expected	52.4%
	Response at Six Months	Expected	23.8%
	Remission at Six Months	Expected	11.1%
	Follow-up at 12 Months	Expected	65.1%
	Response at 12 Months	Above Expected	36.5%
	Remission at 12 Months	Expected	23.8%
Adolescent Mental Health and/or Depression Screening*		Above	98.9%
Colorectal Cancer Screening		Below Expected	60.8%
Optimal Asthma Control - Adults		Expected	46.5%
Optimal Asthma Control - Children		Expected	41.7%
Optimal Diabetes Care		Expected	45.2%
HbA1c Control		Expected	72.1%
BP Control		Expected	84.6%
Daily Aspirin Use		Below Expected	93.3%
Statin Use		Expected	94.2%
Tobacco-Free		Expected	88.5%
Optimal Vascular Care		Expected	43.9%
BP Control		Expected	82.5%
Daily Aspirin Use		Expected	73.7%
Statin Use		Expected	94.7%
Tobacco-Free		Expected	84.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).