

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	89.7%
	Follow-up at Six Months	Expected	42.4%
	Response at Six Months	Expected	11.8%
	Remission at Six Months	Expected	10.6%
	Follow-up at 12 Months	Expected	47.1%
	Response at 12 Months	Expected	14.1%
	Remission at 12 Months	Expected	4.7%
Adult Depression	PHQ-9 Utilization*	Above	89.7%
	Follow-up at Six Months	Expected	47.7%
	Response at Six Months	Expected	15.0%
	Remission at Six Months	Below Expected	5.8%
	Follow-up at 12 Months	Below Expected	44.8%
	Response at 12 Months	Expected	16.5%
	Remission at 12 Months	Below Expected	7.5%
Adolescent Mental Health and/or Depression Screening*		Above	98.8%
Colorectal Cancer Screening		Above Expected	76.4%
Optimal Asthma Control - Adults		Above Expected	72.7%
Optimal Asthma Control - Children		Above Expected	71.1%
Optimal Diabetes Care		Below Expected	41.6%
HbA1c Control		Below Expected	67.9%
BP Control		Above Expected	85.4%
Daily Aspirin Use		Expected	99.0%
Statin Use		Expected	88.5%
Tobacco-Free		Below Expected	78.0%
Optimal Vascular Care		Expected	55.0%
BP Control		Expected	84.5%
Daily Aspirin Use		Above Expected	90.5%
Statin Use		Above Expected	95.8%
Tobacco-Free		Below Expected	73.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).