

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	77.8%
	Follow-up at Six Months	Expected	42.2%
	Response at Six Months	Expected	14.1%
	Remission at Six Months	Expected	10.4%
	Follow-up at 12 Months	Expected	38.5%
	Response at 12 Months	Expected	14.8%
	Remission at 12 Months	Expected	10.4%
Adolescent Mental Health and/or Depression Screening*		Below	77.9%
Colorectal Cancer Screening		Below Expected	62.0%
Optimal Asthma Control - Adults		Below Expected	11.3%
Optimal Asthma Control - Children		Below Expected	3.1%
Optimal Diabetes Care		Below Expected	32.3%
HbA1c Control		Below Expected	62.9%
BP Control		Expected	78.7%
Daily Aspirin Use		Expected	98.3%
Statin Use		Below Expected	78.3%
Tobacco-Free		Expected	81.2%
Optimal Vascular Care		Expected	48.4%
BP Control		Expected	80.6%
Daily Aspirin Use		Expected	84.1%
Statin Use		Below Expected	86.5%
Tobacco-Free		Expected	79.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).