

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	69.2%
	Follow-up at Six Months	Below Expected	18.8%
	Response at Six Months	Expected	9.4%
	Remission at Six Months	Expected	3.1%
	Follow-up at 12 Months	Below Expected	28.1%
	Response at 12 Months	Expected	9.4%
	Remission at 12 Months	Expected	6.3%
Adolescent Mental Health and/or Depression Screening*		Below	81.6%
Colorectal Cancer Screening		Above Expected	73.6%
Optimal Asthma Control - Adults		Below Expected	3.0%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	34.1%
HbA1c Control		Below Expected	64.9%
BP Control		Expected	76.8%
Daily Aspirin Use		Expected	98.5%
Statin Use		Below Expected	77.8%
Tobacco-Free		Expected	80.3%
Optimal Vascular Care		Expected	47.9%
BP Control		Expected	78.4%
Daily Aspirin Use		Expected	83.3%
Statin Use		Below Expected	84.4%
Tobacco-Free		Below Expected	74.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).