

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	87.3%
	Follow-up at Six Months	Expected	51.1%
	Response at Six Months	Expected	13.1%
	Remission at Six Months	Expected	6.8%
	Follow-up at 12 Months	Expected	52.3%
	Response at 12 Months	Expected	16.5%
	Remission at 12 Months	Below Expected	5.1%
Adolescent Mental Health and/or Depression Screening*		Average	93.6%
Colorectal Cancer Screening		Above Expected	83.0%
Optimal Asthma Control - Adults		Expected	43.1%
Optimal Asthma Control - Children		Below Expected	15.5%
Optimal Diabetes Care		Above Expected	52.8%
HbA1c Control		Expected	77.3%
BP Control		Above Expected	87.3%
Daily Aspirin Use		Above Expected	99.8%
Statin Use		Expected	90.4%
Tobacco-Free		Expected	83.6%
Optimal Vascular Care		Expected	60.2%
BP Control		Expected	86.3%
Daily Aspirin Use		Expected	91.5%
Statin Use		Expected	92.9%
Tobacco-Free		Expected	76.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).