

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	92.7%
	Follow-up at Six Months	Expected	34.1%
	Response at Six Months	Expected	15.9%
	Remission at Six Months	Expected	11.4%
	Follow-up at 12 Months	Expected	43.2%
	Response at 12 Months	Expected	15.9%
	Remission at 12 Months	Expected	9.1%
Adolescent Mental Health and/or Depression Screening*		Average	92.2%
Colorectal Cancer Screening		Expected	74.0%
Optimal Asthma Control - Adults		Below Expected	8.7%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	51.6%
HbA1c Control		Expected	80.7%
BP Control		Above Expected	85.9%
Daily Aspirin Use		Expected	99.7%
Statin Use		Expected	86.7%
Tobacco-Free		Expected	83.3%
Optimal Vascular Care		Expected	49.4%
BP Control		Expected	85.5%
Daily Aspirin Use		Expected	88.4%
Statin Use		Expected	90.7%
Tobacco-Free		Below Expected	72.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).