

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	81.7%
	Follow-up at Six Months	Expected	34.7%
	Response at Six Months	Expected	10.2%
	Remission at Six Months	Expected	4.1%
	Follow-up at 12 Months	Expected	32.0%
	Response at 12 Months	Expected	9.5%
	Remission at 12 Months	Expected	3.4%
Adult Depression	PHQ-9 Utilization*	Below	76.9%
	Follow-up at Six Months	Above Expected	58.9%
	Response at Six Months	Expected	21.4%
	Remission at Six Months	Expected	11.2%
	Follow-up at 12 Months	Above Expected	55.2%
	Response at 12 Months	Expected	21.0%
	Remission at 12 Months	Expected	11.2%
Adolescent Mental Health and/or Depression Screening*		Average	92.0%
Colorectal Cancer Screening		Above Expected	74.1%
Optimal Asthma Control - Adults		Expected	50.2%
Optimal Asthma Control - Children		Expected	50.3%
Optimal Diabetes Care		Expected	45.7%
HbA1c Control		Expected	72.7%
BP Control		Expected	78.6%
Daily Aspirin Use		Above Expected	99.3%
Statin Use		Above Expected	92.8%
Tobacco-Free		Expected	82.1%
Optimal Vascular Care		Expected	52.6%
BP Control		Below Expected	75.2%
Daily Aspirin Use		Above Expected	90.9%
Statin Use		Expected	92.9%
Tobacco-Free		Expected	80.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).