

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	75.4%
	Follow-up at Six Months	Expected	47.6%
	Response at Six Months	Expected	16.4%
	Remission at Six Months	Expected	9.5%
	Follow-up at 12 Months	Expected	52.9%
	Response at 12 Months	Expected	20.6%
	Remission at 12 Months	Expected	12.2%
Adolescent Mental Health and/or Depression Screening*		Below	85.3%
Colorectal Cancer Screening		Above Expected	80.9%
Optimal Asthma Control - Adults		Expected	61.9%
Optimal Asthma Control - Children		Expected	53.1%
Optimal Diabetes Care		Above Expected	56.1%
HbA1c Control		Expected	76.1%
BP Control		Expected	85.2%
Daily Aspirin Use		Expected	98.9%
Statin Use		Expected	91.6%
Tobacco-Free		Expected	88.6%
Optimal Vascular Care		Expected	59.0%
BP Control		Expected	83.1%
Daily Aspirin Use		Expected	86.2%
Statin Use		Expected	92.2%
Tobacco-Free		Expected	84.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).