

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	81.8%
	Follow-up at Six Months	Expected	37.4%
	Response at Six Months	Below Expected	6.9%
	Remission at Six Months	Below Expected	2.3%
	Follow-up at 12 Months	Expected	36.2%
	Response at 12 Months	Expected	9.8%
	Remission at 12 Months	Expected	4.6%
Adult Depression	PHQ-9 Utilization*	Average	77.3%
	Follow-up at Six Months	Expected	47.4%
	Response at Six Months	Expected	16.4%
	Remission at Six Months	Expected	9.5%
	Follow-up at 12 Months	Expected	50.2%
	Response at 12 Months	Expected	19.4%
	Remission at 12 Months	Expected	11.2%
Adolescent Mental Health and/or Depression Screening*		Below	87.5%
Colorectal Cancer Screening		Above Expected	76.3%
Optimal Asthma Control - Adults		Expected	46.1%
Optimal Asthma Control - Children		Below Expected	39.6%
Optimal Diabetes Care		Expected	48.4%
HbA1c Control		Expected	72.5%
BP Control		Expected	81.8%
Daily Aspirin Use		Above Expected	99.5%
Statin Use		Above Expected	90.4%
Tobacco-Free		Expected	86.9%
Optimal Vascular Care		Expected	59.3%
BP Control		Expected	81.4%
Daily Aspirin Use		Above Expected	91.5%
Statin Use		Expected	91.2%
Tobacco-Free		Expected	84.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).