

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	79.1%
	Follow-up at Six Months	Expected	73.0%
	Response at Six Months	Expected	21.6%
	Remission at Six Months	Expected	13.5%
	Follow-up at 12 Months	Expected	59.5%
	Response at 12 Months	Expected	18.9%
	Remission at 12 Months	Expected	8.1%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Below Expected	29.3%
Optimal Asthma Control - Adults		Below Expected	25.5%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	35.4%
HbA1c Control		Below Expected	55.5%
BP Control		Expected	76.4%
Daily Aspirin Use		Above Expected	100.0%
Statin Use		Expected	90.9%
Tobacco-Free		Expected	82.3%
Optimal Vascular Care		Expected	54.1%
BP Control		Below Expected	74.1%
Daily Aspirin Use		Above Expected	94.7%
Statin Use		Expected	91.8%
Tobacco-Free		Expected	79.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).