

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Expected	55.3%
	Response at Six Months	Expected	15.8%
	Remission at Six Months	Expected	13.2%
	Follow-up at 12 Months	Expected	50.0%
	Response at 12 Months	Expected	7.9%
	Remission at 12 Months	Expected	7.9%
Adult Depression	PHQ-9 Utilization*	Average	76.9%
	Follow-up at Six Months	Expected	48.2%
	Response at Six Months	Expected	15.7%
	Remission at Six Months	Expected	8.5%
	Follow-up at 12 Months	Expected	49.5%
	Response at 12 Months	Expected	20.3%
	Remission at 12 Months	Expected	10.5%
Adolescent Mental Health and/or Depression Screening*		Average	90.7%
Colorectal Cancer Screening		Above Expected	77.5%
Optimal Asthma Control - Adults		Expected	50.0%
Optimal Asthma Control - Children		Expected	42.2%
Optimal Diabetes Care		Expected	47.4%
HbA1c Control		Expected	75.6%
BP Control		Expected	78.6%
Daily Aspirin Use		Expected	99.5%
Statin Use		Above Expected	91.6%
Tobacco-Free		Expected	83.2%
Optimal Vascular Care		Expected	55.2%
BP Control		Expected	76.1%
Daily Aspirin Use		Above Expected	92.1%
Statin Use		Expected	91.2%
Tobacco-Free		Expected	84.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).