

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	73.1%
	Follow-up at Six Months	Expected	57.5%
	Response at Six Months	Expected	18.3%
	Remission at Six Months	Expected	10.5%
	Follow-up at 12 Months	Expected	52.9%
	Response at 12 Months	Expected	17.7%
	Remission at 12 Months	Expected	11.8%
Adolescent Mental Health and/or Depression Screening*		Average	92.3%
Colorectal Cancer Screening		Below Expected	64.7%
Optimal Asthma Control - Adults		Expected	43.9%
Optimal Asthma Control - Children		Below Expected	22.7%
Optimal Diabetes Care		Expected	46.2%
HbA1c Control		Expected	69.2%
BP Control		Expected	84.1%
Daily Aspirin Use		Expected	99.6%
Statin Use		Above Expected	94.6%
Tobacco-Free		Expected	82.3%
Optimal Vascular Care		Expected	59.5%
BP Control		Expected	80.2%
Daily Aspirin Use		Expected	92.2%
Statin Use		Expected	94.0%
Tobacco-Free		Expected	82.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).